


08/162984



NEW TAXOIDS, THEIR PREPARATION AND
PHARMACEUTICAL COMPOSITIONS CONTAINING THEM

The present invention relates to new taxoids
of general formula:



5 their preparation and pharmaceutical compositions
containing them.

In general formula (I),

Ar represents an aryl radical,

R represents a hydrogen atom or an acetyl, alkoxyacetyl

10 or alkyl radical,

R₁ represents a benzoyl radical or a radical R₂-O-CO- in
which R₂ represents:

- a straight or branched alkyl radical
- containing 1 to 8 carbon atoms, an alkenyl radical
- 15 containing 2 to 8 carbon atoms, an alkynyl radical
- containing 3 to 8 carbon atoms, a cycloalkyl radical
- containing 3 to 6 carbon atoms, a cycloalkenyl radical
- containing 4 to 6 carbon atoms or a bicycloalkyl
- radical containing 7 to 11 carbon atoms, these radicals
- 20 being optionally substituted by one or more
- substituents chosen from halogen atoms and hydroxy
- radicals, alkyloxy radicals containing 1 to 4 carbon

2

atoms, dialkylamino radicals in which each alkyl portion contains 1 to 4 carbon atoms, piperidino radicals, morpholino radicals, 1-piperazinyl radicals (optionally substituted at position 4 by an alkyl radical containing 1 to 4 carbon atoms or by a phenylalkyl radical whose alkyl portion contains 1 to 4 carbon atoms), cycloalkyl radicals containing 3 to 6 carbon atoms, cycloalkenyl radicals containing 4 to 6 carbon atoms, phenyl radicals, cyano radicals, carboxy radicals or alkyloxycarbonyl radicals whose alkyl portion contains 1 to 4 carbon atoms,

- or a phenyl radical optionally substituted by one or more atoms or radicals chosen from halogen atoms and alkyl radicals containing 1 to 4 carbon atoms or alkyloxy radicals containing 1 to 4 carbon atoms,

- or a saturated or unsaturated 4- to 6-membered nitrogen-containing heterocyclyl radical optionally substituted by one or more alkyl radicals containing 1 to 4 carbon atoms, it being understood that the cycloalkyl, cycloalkenyl or bicycloalkyl radicals may be optionally substituted by one or more alkyl radicals containing 1 to 4 carbon atoms.

Preferably, Ar represents a phenyl or α - or β -naphthyl radical optionally substituted by one or more atoms or radicals chosen from halogen atoms (fluorine, chlorine, bromine, or iodine) and alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkoxy, alkylthio,

3

aryloxy, arylthio, hydroxy, hydroxyalkyl, mercapto,
formyl, acyl, acylamino, aroylamino,
alkoxycarbonylamino, amino, alkylamino, dialkylamino,
carboxy, alkoxycarbonyl, carbamoyl, dialkylcarbamoyl,
5 cyano, nitro and trifluoromethyl radicals, it being
understood that the alkyl radicals and the alkyl
portions of the other radicals contain 1 to 4 carbon
atoms, that the alkenyl and alkynyl radicals contain 2
to 8 carbon atoms and that the aryl radicals are phenyl
10 or α - or β -naphthyl radicals or alternatively Ar
represents a 5-membered aromatic heterocyclic radical
containing one or more atoms, which are identical or
different, chosen from nitrogen, oxygen or sulphur
atoms, optionally substituted by one or more
15 substituents, which are identical or different, chosen
from halogen atoms (fluorine, chlorine, bromine or
iodine) and alkyl radicals containing 1 to 4 carbon
atoms, aryl radicals containing 6 to 10 carbon atoms,
alkoxy radicals containing 1 to 4 carbon atoms, aryloxy
20 radicals containing 6 to 10 carbon atoms, amino
radicals, alkylamino radicals containing 1 to 4 carbon
atoms, dialkylamino radicals in which each alkyl
portion contains 1 to 4 carbon atoms, acylamino
radicals in which the acyl portion contains 1 to 4
25 carbon atoms, alkoxycarbonylamino radicals containing 1
to 4 carbon atoms, acyl radicals containing 1 to 4
carbon atoms, arylcarbonyl radicals in which the aryl
portion contains 6 to 10 carbon atoms, cyano radicals,

4

carboxy radicals, carbamoyl radicals, alkylcarbamoyl radicals in which the alkyl portion contains 1 to 4 carbon atoms, dialkylcarbamoyl radicals in which each alkyl portion contains 1 to 4 carbon atoms or
5 alkoxy carbonyl radicals in which the alkoxy portion contains 1 to 4 carbon atoms.


More particularly, Ar represents a phenyl, 2- or 3-thienyl or 2- or 3-furyl radical optionally substituted by one or more atoms or radicals, which are
10 identical or different, chosen from halogen atoms and alkyl, alkoxy, amino, alkylamino, dialkylamino, acylamino, alkoxy carbonylamino and trifluoromethyl radicals.

Still more particularly, Ar represents a
15 phenyl radical optionally substituted by a chlorine or fluorine atom or by an alkyl (methyl), alkoxy (methoxy), dialkylamino (diethylamino), acylamino (acetyl amino) or alkoxy carbonylamino (tert-butoxy carbonylamino) or 2- or 3-thienyl or 2- or
20 3-furyl radical.

Of even more special interest are the products of general formula (I) in which Ar represents a phenyl radical and R₁ represents a benzoyl or tert-butoxy carbonyl radical.

25 According to the present invention, the new taxoids of general formula (I) can be obtained from a product of general formula:

5




in which Ar and R₁ are defined as above and R₂ and R₄, which are identical or different represent a hydrogen atom or an alkyl radical containing 1 to 4 carbon atoms, or an aralkyl radical whose alkyl portion contains 1 to 4 carbon atoms and the aryl portion preferably represents a phenyl radical optionally substituted by one or more alkoxy radicals containing 1 to 4 carbon atoms, or an aryl radical preferably representing a phenyl radical optionally substituted by one or more alkoxy radicals containing 1 to 4 carbon atoms, or alternatively R₂ represents an alkoxy radical containing 1 to 4 carbon atoms or a trihalomethyl radical such as trichloromethyl or a phenyl radical substituted by a trihalomethyl radical such as trichloromethyl and R₄ represents a hydrogen atom, or alternatively R₂ and R₄ form, together with the carbon atom to which they are attached, a 4- to 7-membered ring, and G₁ represents a hydrogen atom or an acetyl, alkoxyacetyl or alkyl radical or a hydroxy-protecting group, the procedure being carried out, according to the meanings of R₂ and R₄, in the following manner:

1) when R₂ represents a hydrogen atom or an

6

alkoxy radical containing 1 to 4 carbon atoms or an optionally substituted aryl radical and R_1 represents a hydrogen atom, the product of general formula (II) is treated in acidic medium in order to obtain a product of general formula:



in which Ar , R_1 and G_1 are defined as above, whose G_1 radical is, if necessary, replaced by a hydrogen atom.

The deprotection of the side chain of the product of general formula (II) can also be carried out in the presence of an inorganic acid (hydrochloric acid or sulphuric acid) or an organic acid (acetic acid, methanesulphonic acid, trifluoromethanesulphonic acid or p-toluenesulphonic acid), used alone or in the form of a mixture, the procedure being carried out in an organic solvent chosen from alcohols (methanol, ethanol or isopropanol), ethers (tetrahydrofuran, diisopropyl ether or methyl t-butyl ether), esters (ethyl acetate, isopropyl acetate or n-butyl acetate), aliphatic hydrocarbons (pentane, hexane or heptane), halogenated aliphatic hydrocarbons (dichloromethane or 1,2-dichloroethane), aromatic hydrocarbons (benzene, toluene or xylenes) and nitriles (acetonitrile) at a

temperature of between -10 and 60°C, preferably between 15 and 30°C. The acid may be used in a catalytic or stoichiometric quantity or in excess.

The deprotection can also be carried out
5 under oxidizing conditions, using for example ammonium cerium(IV) nitrate in an acetonitrile-water mixture or 2,3-dichloro-5,6-dicyano-1,4-benzoquinone in water.

The deprotection can also be carried out
under reducing conditions, for example by
10 hydrogenolysis in the presence of a catalyst.


When G₁ represents a protecting group, it is preferably a 2,2,2-trichloroethoxycarbonyl or 2-(2-trichloromethylpropoxy)carbonyl radical whose replacement by a hydrogen atom is carried out using
15 zinc, optionally combined with copper, in the presence of acetic acid, at a temperature of between 20 and 60°C or by means of an inorganic or organic acid such as hydrochloric acid or acetic acid in a solution in an aliphatic alcohol containing 1 to 3 carbon atoms or in
20 an aliphatic ester such as ethyl acetate, isopropyl acetate or n-butyl acetate in the presence of zinc optionally combined with copper, or alternatively, when G₁ represents an alkoxycarbonyl radical, its optional replacement by a hydrogen atom is carried out by
25 treatment in alkaline medium or by the action of a zinc halide under conditions which do not affect the rest of the molecule. Generally, the alkaline treatment is carried out by the action of ammonia in aqueous-

8

alcoholic medium, at a temperature close to 20°C.

Generally, the treatment with a zinc halide, preferably zinc iodide is carried out in methanol at a temperature close to 20°C.

- 5 2) when R_1 and R_4 , which are identical or different, represent an alkyl radical containing 1 to 4 carbon atoms, or an aralkyl radical whose alkyl portion contains 1 to 4 carbon atoms and the aryl portion is preferably an optionally substituted phenyl radical, or
10 alternatively R_1 represents a trihalomethyl radical or a phenyl radical substituted by a trihalomethyl radical and R_4 represents a hydrogen atom, or alternatively R_1 and R_4 form, together with the carbon atom to which they are attached, a 4- to 7-membered ring, the product of
15 general formula (II) is converted to the product of general formula:



in which Ar and G_1 are defined as above, which is acylated by means of benzoyl chloride or a reactive derivative of general formula:



in which R_2 is defined as above and X represents a halogen atom (fluorine or chlorine) or a residue $-O-R_2$

or $-O-CO-O-R_2$, to give a product of general formula (III) in which Ar, R_1 and G_1 are defined as above, whose G_1 radical is, if necessary, replaced by a hydrogen atom.


5 The products of general formula (IV) can be obtained by treating a product of general formula (II), in which Ar, R_1 and G_1 are defined as above, R_2 and R_4 , which are identical or different, represent an alkyl, aralkyl or aryl radical, or alternatively R_2 and R_4 form
10 together with the carbon atom to which they are attached a 4- to 7-membered ring, with an inorganic acid (hydrochloric acid or sulphuric acid) or an organic acid (formic acid) optionally in an alcohol containing 1 to 3 carbon atoms (methanol, ethanol or
15 isopropanol) at a temperature of between 0 and 50°C. Preferably, formic acid is used at a temperature close to 20°C.

 The acylation of the product of general formula (IV) by means of benzoyl chloride or a reactive
20 derivative of general formula (V) is carried out in an inert organic solvent chosen from esters such as ethyl acetate, isopropyl acetate or n-butyl acetate and halogenated aliphatic hydrocarbons such as dichloromethane or 1,2-dichloroethane in the presence
25 of an inorganic base such as sodium bicarbonate or an organic base such as triethylamine. The reaction is carried out at a temperature of between 0 and 50°C, preferably close to 20°C.


10

When the radical G_1 represents a protecting group, its replacement by a hydrogen atom is carried out under the conditions described above.

The products of general formula (II) can be obtained according to one of the following methods:
 5 1) by esterification of the product of general formula:



in which G_1 is defined as above, by means of an acid of general formula:



in which Ar , R_1 , R_3 and R_4 are defined as above, or of a
 10 derivative of this acid.

The esterification by means of an acid of general formula (VII) can be carried out in the presence of a condensing agent (carbodiimide, reactive carbonate) and an activating agent (aminopyridine) in
 15 an organic solvent (ether, ester, ketones, nitriles, aliphatic hydrocarbons, halogenated aliphatic hydrocarbons or aromatic hydrocarbons) at a temperature


11

of between -10 and 90°C.

The esterification may also be performed using the acid of general formula (VII) in anhydride form, the procedure being carried out in the presence of an activating agent (aminopyridine) in an organic solvent (ethers, esters, ketones, nitriles, aliphatic hydrocarbons, halogenated aliphatic hydrocarbons or aromatic hydrocarbons) at a temperature of between 0 and 90°C.

The esterification can also be performed using the acid of general formula (VII) in halide form or in anhydride form with an aliphatic or aromatic acid, optionally prepared in situ, in the presence of a base (tertiary aliphatic amine), the procedure being carried out in an organic solvent (ethers, esters, ketones, nitriles, aliphatic hydrocarbons, halogenated aliphatic hydrocarbons or aromatic hydrocarbons) at a temperature of between 0 and 80°C.

The acid of general formula (VII) can be obtained by saponification of an ester of general formula:



in which Ar, R₁, R₃ and R₄ are defined as above and R₅ represents an alkyl radical containing 1 to 4 carbon atoms optionally substituted by a phenyl radical.


12

Generally, the saponification is carried out by means of an inorganic base (alkali metal hydroxide, carbonate or bicarbonate) in aqueous-alcoholic medium (methanol-water) at a temperature of between 10 and 40°C.

The ester of general formula (VIII) can be obtained by the action of a product of general formula:




in which R_3 and R_4 are defined as above in the form of a dialkylacetal or an enol alkyl ether, on an ester of general formula:



in which Ar , R_1 and R_5 are defined as above, the procedure being carried out in an inert organic solvent (aromatic hydrocarbon) in the presence of a strong inorganic acid (sulphuric acid) or organic acid (p-toluenesulphonic acid optionally in the form of a pyridinium salt) at a temperature of between 0°C and the boiling temperature of the reaction mixture.


The ester of general formula (X) can be obtained by the action of a product of general formula (V) on an ester of general formula:

13



in which Ar and R₅ are defined as above, the procedure being carried out in an organic solvent (ester, halogenated aliphatic hydrocarbon) in the presence of an inorganic or organic base at a temperature of
5 between 0 and 50°C.


The product of general formula (XI) can be obtained by reduction of an azide of general formula:



in which Ar and R₅ are defined as above, by means of hydrogen in the presence of a catalyst such as
10 palladium on carbon, the procedure being carried out in an organic solvent (ester).


The product of general formula (XII) can be obtained by the action of an azide such as trimethylsilyl azide in the presence of zinc chloride
15 or an alkali metal (sodium, potassium or lithium) azide in aqueous-organic medium (water-tetrahydrofuran) at a temperature of between 20°C and the boiling temperature of the reaction mixture, on an epoxide of general formula:

14



in which Ar and R₅ are defined as above, optionally prepared in situ.

The epoxide of general formula (XIII) can be obtained, optionally in situ, by dehydrohalogenation of
5 a product of general formula:



in which Ar is defined as above, Hal represents a halogen atom, preferably a bromine atom, and R₆ and R₇, which are identical or different, represent a hydrogen atom or an alkyl radical containing 1 to 4 carbon atoms
10 or a phenyl radical, at least one being an alkyl radical or a phenyl radical, by means of an alkali-metal alcoholate, optionally prepared in situ, in an inert organic solvent such as tetrahydrofuran at a temperature of between -80°C and 25°C.


15 The product of general formula (XIV) can be obtained by the action of an aldehyde of general formula:



in which Ar is defined as above, on a halide of general

15

formula:




in which Hal, R₆ and R₇ are defined as above, anionized beforehand.

Generally, the procedure is carried out in an inert organic solvent chosen from ethers (ethyl ether) and halogenated aliphatic hydrocarbons (methylene chloride) at a temperature of between -80 and 25°C, in the presence of a tertiary amine (triethylamine) and an enolysing agent (di-n-butylboron triflate).

The product of general formula (XVI) can be obtained by the action of a halide of a haloacetic acid, preferably bromoacetic acid bromide, on the corresponding oxazolidinone.

The product of general formula (XI) can be obtained by hydrogenolysis of a product of general formula:



in which Ar and R₅ are defined as above and Ph represents an optionally substituted phenyl radical.


16

Generally, the hydrogenolysis is carried out by means of hydrogen in the presence of a catalyst. More particularly, palladium on carbon containing 1 to 10 % by weight of palladium or palladium dihydroxide
5 containing 20 % by weight of palladium is used as catalyst.

The hydrogenolysis is carried out in an organic solvent or in a mixture of organic solvents. It is advantageous to carry out the procedure in acetic
10 acid optionally combined with an aliphatic alcohol containing 1 to 4 carbon atoms such as a mixture of acetic acid-methanol at a temperature of between 20 and 80°C.

The hydrogen necessary for the hydrogenolysis
15 can also be provided by a compound which liberates hydrogen by chemical reaction or by thermal decomposition (ammonium formate). It is advantageous to carry out the procedure at a hydrogen pressure of between 1 and 50 bar.

20 The product of general formula (XVII) can be obtained by hydrolysis or alcoholysis of a product of general formula:




in which Ar and Ph are defined as above.

17

It is particularly advantageous to carry out an alcoholysis by means of an alcohol of formula R_3 -OH in which R_3 is defined as above, the procedure being carried out in acidic medium.

5 Preferably, the alcoholysis is carried out by means of methanol in the presence of a strong inorganic acid such as hydrochloric acid at a temperature close to the reflux temperature of the reaction mixture.

The product of general formula (XVIII) can be
10 obtained by saponification of an ester of general formula:



in which Ar and Ph are defined as above and R_8 represents an alkyl, phenylalkyl or phenyl radical, followed by separation of the 3R, 4S diastereoisomer of
15 general formula (XVII) from the other diastereoisomers.


Generally, the saponification is carried out by means of an inorganic or organic base such as ammonium hydroxide, lithium hydroxide, sodium hydroxide or potassium hydroxide in a suitable solvent such as a
20 methanol-water or tetrahydrofuran-water mixture at a temperature of between -10°C and 20°C .

The separation of the 3R,4S diastereoisomer can be carried out by selective crystallization from a


18

suitable organic solvent such as ethyl acetate.

The product of general formula (XIX) can be obtained by cycloaddition of an imine of general formula:



5 in which Ar and Ph are defined as above, onto an acid halide of general formula:



in which R₈ is defined as above and Y represents a halogen atom such as a bromine or chlorine atom.


Generally, the reaction is carried out at a
10 temperature of between 0 and 50°C in the presence of a base chosen from aliphatic tertiary amines (triethylamine) or pyridine in an organic solvent chosen from optionally halogenated aliphatic hydrocarbons (methylene chloride or chloroform) and
15 aromatic hydrocarbons (benzene, toluene or xylenes).

The product of general formula (XX) can be obtained under conditions analogous to those described by M. Furukawa et al., Chem. Pharm. Bull., 25 (1), 181-184 (1977).

20 The product of general formula (VI) can be obtained by the action of an alkali metal halide

19

(sodium iodide or potassium fluoride) or an alkali metal azide (sodium azide) or a quaternary ammonium salt or an alkali metal phosphate, on a baccatin III or 10-deacetylbaccatin III derivative of general formula:



5 in which G_1 is defined as above.

Generally, the reaction is carried out in an organic solvent chosen from ethers (tetrahydrofuran, diisopropyl ether, methyl t-butyl ether) and nitriles (acetonitrile), alone or in the form of a mixture, at a
10 temperature of between 20°C and the boiling temperature of the reaction mixture.

The product of formula (XXII) in which G_1 represents a hydrogen atom or an acetyl, alkoxyacetyl or alkyl radical can be obtained by the action of a
15 trifluoromethanesulphonic acid derivative such as the anhydride or N-phenyltrifluoromethanesulphonimide, on baccatin III or 10-deacetylbaccatin III, which can be extracted according to known methods from yew leaves (*Taxus baccata*), optionally followed by protection in
20 position 10, it being understood that in order to obtain a product of general formula (XXII) in which G_1 represents an alkoxyacetyl or alkyl radical, it is

20

necessary to treat beforehand the 10-deacetylbaccatin III protected in position 7, preferably with a silylated radical, with an alkoxy acetic acid halide or with an alkyl halide.


5 Generally, the reaction of a trifluoromethanesulphonic acid derivative is carried out in an inert organic solvent (optionally halogenated aliphatic hydrocarbons, or aromatic hydrocarbons) in the presence of an organic base such as an aliphatic
10 tertiary amine (triethylamine) or pyridine, at a temperature of between -50 and +20°C.

 Generally, the introduction of an alkoxyacetyl group is carried out by treating the protected 10-deacetylbaccatin III with an alkoxyacetic
15 acid halide, the procedure being carried out in a basic organic solvent such as pyridine at a temperature close to 20°C.

 Generally, the introduction of an alkyl radical is carried out by treating the
20 10-deacetylbaccatin III, protected and metallized in position 10, by means, for example, of an alkali metal hydride (sodium hydride) or a metallic alkylide (butyllithium), with an alkyl halide.

2) by the action of an alkali metal halide (sodium
25 iodide or potassium fluoride) or an alkali metal azide (sodium azide) or a quaternary ammonium salt or an alkali metal phosphate on a product of general formula:


21



in which Ar, R₁, R₃, R₄ and G₁ are defined as above.

Generally, the reaction is carried out in an organic solvent chosen from ethers (tetrahydrofuran, diisopropyl ether or methyl t-butyl ether) and nitriles (acetonitrile), alone or in the form of a mixture, at a temperature of between 20°C and the boiling temperature of the reaction mixture.

The product of general formula (XXIII) can be obtained by the action of a trifluoromethanesulphonic acid derivative such as the anhydride or N-phenyltrifluoromethanesulphonimide on a taxoid of general formula:




in which Ar, R₁, R₃, R₄ and G₁ are defined as above.

Generally, the reaction is carried out in an inert organic solvent (optionally halogenated aliphatic hydrocarbons, or aromatic hydrocarbons) in the presence

22

of an organic base such as an aliphatic tertiary amine (triethylamine) or pyridine, at a temperature of between -50 and +20°C.

The taxoid of general formula (XXIV), in which G_1 represents a hydrogen atom or an acetyl radical, can be obtained from a product of general formula:



in which Ar, R_1 , R_3 and R_4 are defined as above, G'_1 represents a hydroxy-protecting group and G'_2 represents an acetyl, alkoxyacetyl or alkyl radical or a hydroxy-protecting group, by replacement of the protecting groups G'_1 and optionally G'_2 by hydrogen atoms.

The radicals G'_1 and G'_2 , when they represent a hydroxy-protecting group, are preferably 2,2,2-trichloroethoxycarbonyl or 2-(2-trichloromethyl-propoxy)carbonyl radicals or trialkylsilyl, dialkylarylsilyl, alkyl diarylsilyl or triarylsilyl radicals in which the alkyl portions contain 1 to 4 carbon atoms and the aryl portions are preferably phenyl radicals, it being possible, in addition, for G'_2 to represent an alkoxyacetyl radical.

When G'_1 and G'_2 represent a 2,2,2-

23

trichloroethoxycarbonyl or 2-(2-trichloromethylpro-
poxy)carbonyl radical, the replacement of the
protecting groups by hydrogen atoms is carried out
using zinc, optionally combined with copper, in the
5 presence of acetic acid at a temperature of between 20
and 60°C or by means of an inorganic or organic acid
such as hydrochloric acid or acetic acid in solution in
an aliphatic alcohol containing 1 to 3 carbon atoms or
an aliphatic ester such as ethyl acetate, isopropyl
10 acetate or n-butyl acetate in the presence of zinc
optionally combined with copper.

When G', represents a silylated radical and
G', represents an acetyl, alkoxyacetyl or alkyl radical,
the replacement of the protecting group G', by a
15 hydrogen atom can be carried out by means of, for
example, gaseous hydrochloric acid in ethanolic
solution at a temperature close to 0°C, under
conditions which are without effect on the rest of the
molecule.


20 When G', represents an alkoxyacetyl radical,
its optional replacement by a hydrogen atom is carried
out by treatment in alkaline medium or by the action of
a zinc halide under conditions which do not affect the
rest of the molecule. Generally, the alkaline treatment
25 is carried out by the action of ammonia in aqueous-
alcoholic medium, at a temperature close to 20°C.
Generally, the treatment with a zinc halide, preferably
zinc iodide, is carried out in methanol at a

24

temperature close to 20°C.


The product of general formula (XXV) can be obtained under the conditions described in international application PCT/WO 9209589.

5 The new derivatives of general formula (I) can also be obtained by esterification of a product of general formula (VI) by means of an acid of general formula:



in which Ar and R₁ are defined as above and G₃,
10 represents a hydroxy-protecting group chosen from methoxymethyl, 1-ethoxyethyl, benzyloxymethyl, (β-trimethylsilyloxy)methyl, tetrahydropyranyl, 2,2,2-trichloroethoxymethyl, 2,2,2-trichloroethoxycarbonyl or 2-(2-
15 trichloromethylpropoxy)carbonyl radicals or CH₂-Ph radicals in which Ph represents a phenyl radical optionally substituted by one or more atoms or radicals, which are identical or different, chosen from halogen atoms and alkyl radicals containing 1 to 4
20 carbon atoms or alkoxy radicals containing 1 to 4 carbon atoms, or an activated derivative of this acid, to give a product of general formula:

25



in which Ar, R_1 , G_1 , G_2 and G_3 are defined as above, followed by the replacement of the protecting groups G_1 , G_2 and G_3 by hydrogen atoms to give a product of general formula (I).


5 The esterification can be performed under the conditions described above for the esterification of the product of general formula (VI) by means of an acid of general formula (VII).

 The replacement of the protecting groups G_1 and G_2 of the product of general formula (XXVII) by a
 10 hydrogen atom is carried out by treatment with zinc, optionally combined with copper, in the presence of acetic acid at a temperature of between 30 and 60°C or by means of an inorganic or organic acid such as
 15 hydrochloric acid or acetic acid in solution in an aliphatic alcohol containing 1 to 3 carbon atoms or an aliphatic ester such as ethyl acetate, isopropyl acetate or n-butyl acetate in the presence of zinc optionally combined with copper, when G_1 and G_2
 20 represent a 2,2,2-trichloroethoxycarbonyl or 2-(2-trichloromethylpropoxy)carbonyl radical. The

26

replacement of the protecting group G_1 , when it represents a silylated radical or an acetal residue, can be carried out by treatment in acidic medium such as for example hydrochloric acid in solution in an aliphatic alcohol containing 1 to 3 carbon atoms (methanol, ethanol, propanol or isopropanol) or aqueous hydrofluoric acid at a temperature of between 0 and 40°C, when it represents an acetal residue, the replacement of the protecting group G_1 then being carried out under the conditions described above. When G_1 represents a group $-\text{CH}_2-\text{Ph}$, the replacement of this protecting group with a hydrogen atom can be carried out by hydrogenolysis in the presence of a catalyst.

The acid of general formula (XXVI) can be obtained by saponification of an ester of general formula:



in which Ar, R_1 , R_5 , and G_3 are defined as above.

Generally, the saponification is carried out by means of an inorganic base (alkali metal hydroxide, carbonate or bicarbonate) in aqueous-alcoholic medium (methanol-water) at a temperature of between 10 and 40°C.

The ester of general formula (XXVIII) can be obtained according to the usual methods for the preparation of ethers, and more particularly according

to the procedures described by J-N. DENIS et al.,
J. Org. Chem., 51, 46-50 (1986), from a product of
general formula (XI).

The new products of general formula (I)
5 obtained using the procedures according to the
invention can be purified according to known methods
such as crystallization or chromatography.

The products of general formula (I) have
remarkable biological properties.

10 In vitro, measurement of the biological
activity is carried out on tubulin extracted from pig
brain by the method of M.L. Shelanski et al. Proc.
Natl. Acad. Sci. USA, 70, 765-768 (1973). The study of
the depolymerization of the microtubules into tubulin
15 is carried out according to the method of G. Chauvière
et al., C.R. Acad. Sci., 293, série II, 501-503 (1981).
In this study, the products of general formula (I)
proved at least as active as taxol and Taxotere.

In vivo, the products of general formula (I)
20 proved active in mice grafted with the B16 melanoma at
doses of between 1 and 10 mg/kg intraperitoneally, as
well as on other liquid or solid tumours.

The new compounds have anti-tumor properties, more
particularly, activity against tumors which are resistant to
25 Taxol® and Taxotere®. Such tumors include, for example, colon
tumors which have an elevated expression of *mdr 1* gene (multi-

drug resistant gene). Multi-drug resistance is the usual term relating to the resistance by a tumor against various compounds having differing structures and mechanisms of action. Taxoids are generally known to be highly recognized
5 by experimental tumors such as P388/DOX, a P388 murine leukemia cell line selected for doxorubicin (DOX) resistance, which express mdr 1. The new compounds according to the present invention are less recognized by P388/DOX. More particularly, the new compounds are less recognized than
10 Taxotere® by mdr 1.

In particular, it has been found that the new compounds of the present invention including the compounds of example 1, example 2 and example 3 have better multi-drug
15 resistance properties than Taxol® and Taxotere®. Additionally it has surprisingly been found that the compound of example 3 has substantially better multi-drug resistance properties than the compounds of example 1 and example 2.

20 The following examples illustrate the present invention.

EXAMPLE 1

25 A solution of 2.01 g of 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,10 β -dihydroxy-7 β ,8 β -

29

methylene-9-oxo-19-nor-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinecarboxylate in 20 cm³ of formic acid is stirred for 4 hours at a temperature close to 20°C and then concentrated to dryness under reduced pressure (0.27 kPa) at 40°C. The foam obtained is dissolved in 100 cm³ of dichloromethane and the solution obtained is supplemented with 20 cm³ of a saturated aqueous sodium hydrogen carbonate solution. The aqueous phase is separated after settling has taken place and extracted with 20 cm³ of dichloromethane. The organic phases are pooled, dried over magnesium sulphate, filtered and then concentrated to dryness under reduced pressure (2.7 kPa) at 40°C. 1.95 g of a white foam are obtained which are purified by chromatography on 200 g of silica (0.063-0.2 mm) contained in a column 7 cm in diameter, eluting with a dichloromethane-methanol mixture (98-2 by volume) and collecting 30 cm³ fractions. The fractions containing only the desired product are pooled and concentrated to dryness under reduced pressure (0.27 kPa) at 40°C for 2 hours. 1.57 g of 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,10 β -dihydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (2R,3S)-3-amino-2-hydroxy-3-phenylpropionate are obtained in the form of a white foam.

To a solution of 400 mg of 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,10 β -dihydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (2R,3S)-3-amino-

2-hydroxy-3-phenylpropionate in 1 cm³ of dichloromethane, kept under an argon atmosphere, are added 60 mg of sodium hydrogen carbonate and then, dropwise, at a temperature close to 20°C, a solution of 5 0.16 g of di-tert-butyl dicarbonate in 1 cm³ of dichloromethane. The solution obtained is stirred for 64 hours at a temperature close to 20°C and then supplemented with a mixture of 5 cm³ of distilled water and 10 cm³ of dichloromethane. The organic phase is 10 washed with three times 2 cm³ of distilled water. The organic phase is dried over magnesium sulphate, filtered and then concentrated to dryness under reduced pressure (2.7 kPa) at 40°C. 317 mg of a white foam are thus obtained which are purified by chromatography on 15 30 g of silica (0.063-0.2 mm) contained in a column 3 cm in diameter, eluting with a dichloromethane-methanol mixture (95-5 by volume) and collecting 5 cm³ fractions. The fractions containing only the desired product are pooled and concentrated to dryness under 20 reduced pressure (0.27 kPa) at 40°C for 2 hours. 161 mg of 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,10 β -dihydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (2R,3S)-3-tert-butoxycarbonylamino-2-hydroxy-3-phenylpropionate are thus obtained in the form of a 25 white foam whose characteristics are the following:

- specific rotation: $[\alpha]_D^{20} = -17^\circ$ (c = 0.482; methanol)
- proton NMR spectrum: (400 MHz; CDCl₃; temperature of 323 K; δ in ppm; coupling constants J in Hz): 1.21 (s,

31

3H:-CH₃, 16 or 17); 1.28 (s, 3H:-CH₃, 16 or 17); 1.34 [s, 9H:-C(CH₃)₃]; from 1.30 to 1.50 (mt, 1H:-H₇); 1.80 and 2.36 (2mt, 1H each:-CH₂- of cyclopropane); 1.88 (s, 3H:-CH₃, 18); 2.13 [mt, 1H:-(CH)-H₆]; 2.26 [dd, 1H, J = 15 to 8.5:-(CH)-H₁₄]; 2.35 (s, 3H:-COCH₃); from 2.35 to 2.50 [mt, 2H:-(CH)-H₁₄ and -(CH)-H₆]; 3.21 (d, 1H, J = 4:-OH_{2'}); 4.08 [d, 1H, J = 8:-(CH)-H₂₀]; 4.16 (d, 1H, J = 7:-H₃); 4.18 (s, 1H, -OH₁₀); 4.31 [d, 1H, J = 8:-(CH)-H₂₀]; 4.61 (dd, 1H, J = 4 and 2:-H_{2'}); 4.74 (d, 1H, J = 4:-H₅); 5.00 (s, 1H:-H₁₀); 5.26 (dd, 1H, J = 9 and 2:-H_{3'}); 5.33 (d, 1H, J = 9:-NH_{3'}); 5.69 (d, 1H, J = 7:-H₂); 6.29 (d, 1H, J = 8.5:-H₁₃); from 7.30 to 7.50 [mt, 5H:-C₆H₅ in 3' (-H₂ to -H₆); 7.51 [t, 2H, J = 7.5:-OCOC₂H₅ (-H₃ and H₅)]; 7.60 [t, 1H, J = 7.5:-OCOC₂H₅ (-H₄)]; 8.14 [d, 2H, J = 7.5:-OCOC₂H₅ (-H₂ and H₆)].

The 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,10 β -dihydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinecarboxylate can be prepared in the following manner:

To a solution of 2.5 g of 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,10 β -dihydroxy-9-oxo-7 β -trifluoromethanesulphonate-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinecarboxylate in 25 cm³ of anhydrous acetonitrile and 3 cm³ of anhydrous tetrahydrofuran, kept under an argon atmosphere, are added 2.5 g of

32

- sodium azide. The reaction mixture is heated for 2 hours, with stirring and under an argon atmosphere at a temperature close to 80°C, then cooled to a temperature close to 20°C and supplemented with 30 cm³ of distilled water. The aqueous phase is separated by decantation and then extracted with 20 cm³ of dichloromethane. The combined organic phases are dried over magnesium sulphate, filtered and then concentrated to dryness under reduced pressure (2.7 kPa) at 40°C.
- 2.44 g of a yellow foam are thus obtained which are purified by chromatography on 300 g of silica (0.063-0.2 mm) contained in a column 8 cm in diameter, eluting with a dichloromethane-ethyl acetate mixture (90-10 by volume) and collecting 60 cm³ fractions.
- Fractions 47 to 70 are pooled and concentrated to dryness under reduced pressure (0.27 kPa) at 40°C for 2 hours. 2.01 g of 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,10 β -dihydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinecarboxylate are thus obtained in the form of a white foam.

The 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,10 β -dihydroxy-9-oxo-7 β -trifluoromethanesulphonate-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinecarboxylate can be prepared in the following manner:

To a solution of 2.86 g of 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,7 β ,10 β -trihydroxy-9-oxo-11-

33

taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinecarboxylate in 29 cm³ of anhydrous dichloromethane, kept under an argon atmosphere, are added 0.955 cm³ of pyridine and 50 mg of powdered activated 4Å molecular sieve. The reaction mixture is cooled to a temperature close to -35°C, slowly supplemented with 0.85 cm³ of trifluoromethanesulphonic anhydride, stirred at a temperature close to -5°C for 15 minutes and supplemented with 10 cm³ of distilled water. After filtration on sintered glass provided with celite and rinsing off the sintered glass with 3 times 10 cm³ of a methanol-dichloromethane mixture (10-90 by volume), the aqueous phase is separated after settling has taken place and extracted with twice 10 cm³ of dichloromethane. The organic phases are pooled, dried over magnesium sulphate, filtered and then concentrated to dryness under reduced pressure (2.7 kPa) at 40°C. 3.87 g of a white foam are obtained which are purified by chromatography on 400 g of silica (0.063-0.2 mm) contained in a column 10 cm in diameter, eluting with a dichloromethane-ethyl acetate gradient (from 97.5-2.5 to 90-10 by volume) and collecting 80 cm³ fractions. The fractions containing only the desired product are pooled and concentrated to dryness under reduced pressure (0.27 kPa) at 40°C for 2 hours. 3.0 g of 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,10 β -dihydroxy-9-oxo-7 β -trifluoromethanesulphonate-11-taxen-13 α -yl

(4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinecarboxylate are thus obtained in the form of a white foam.

The 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-
5 1 β ,7 β ,10 β -trihydroxy-9-oxo-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinecarboxylate can be prepared in the following manner:

A solution of 24.35 g of 4-acetoxy-2 α -
10 benzoyloxy-5 β ,20-epoxy-9-oxo-7 β ,10 β -[bis(2,2,2-trichloroethoxy)carbonyloxy]-1 β -hydroxy-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinecarboxylate in a mixture of 130 cm³ of ethyl acetate and 46.5 cm³ of acetic acid is heated, with
15 stirring and under an argon atmosphere up to a temperature close to 60°C and then supplemented with 40 g of zinc powder. The reaction mixture is then stirred for 30 minutes at 60°C and then cooled to a temperature close to 20°C and filtered on sintered
20 glass provided with celite. The sintered glass is washed with 100 cm³ of a methanol-dichloromethane mixture (20-80 by volume); the filtrates are pooled and then concentrated to dryness under reduced pressure (0.27 kPa) at a temperature close to 40°C.

25 The residue is supplemented with 500 cm³ of dichloromethane. The organic phase is washed with twice 50 cm³ of a saturated aqueous sodium hydrogen carbonate solution and then with 50 cm³ of distilled water. The

35

aqueous phases obtained after settling has taken place and pooled are extracted with twice 30 cm³ of dichloromethane. The organic phases are pooled, dried over magnesium sulphate, filtered and then concentrated to dryness under reduced pressure (2.7 kPa) at 40°C. 19.7 g of a white foam are obtained which are purified by chromatography on 800 g of silica (0.063-0.2 mm) contained in a column 10 cm in diameter, eluting with a dichloromethane-methanol gradient (from 100-0 to 97-3 by volume) and collecting 80 cm³ fractions. The fractions containing only the desired product are pooled and concentrated to dryness under reduced pressure (0.27 kPa) at 40°C for 2 hours. 16.53 g of 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,7 β ,10 β -trihydroxy-9-oxo-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinecarboxylate in the form of a white foam.

The 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-9-oxo-7 β ,10 β -[bis(2,2,2-trichloroethoxy)carbonyloxy]-1 β -hydroxy-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinecarboxylate can be prepared according to the method described in international application PCT WO 9209589.

EXAMPLE 2

To a solution of 550 mg of 4 α ,10 β -diacetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β -hydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (2R,3S)-3-amino-2-hydroxy-3-

36

phenylpropionate are added 45 cm³ of distilled water, 45 cm³ of a saturated aqueous sodium hydrogen carbonate solution and then, dropwise, at a temperature close to 20°C, 0.096 cm³ of benzoyl chloride. The mixture

5 obtained is stirred for 10 minutes at a temperature close to 20°C. After settling has taken place, the aqueous phase is extracted with twice 30 cm³ of ethyl acetate. The combined organic phases are dried over magnesium sulphate, filtered and then concentrated to

10 dryness under reduced pressure (2.7 kPa) at 40°C. 670 mg of a white foam are thus obtained which are purified by chromatography at atmospheric pressure on 50 g of silica (0.063-0.2 mm) contained in a column 2.5 cm in diameter, eluting with a methanol-

15 dichloromethane mixture (1-99 then 2.5-97.5 by volume) and collecting 10 cm³ fractions. The fractions containing only the desired product are pooled and concentrated to dryness under reduced pressure (2.7 kPa) at 40°C. 610 mg of a white foam are thus

20 obtained. A sample of 300 mg is purified by preparative chromatography on 12 thin-layer silica plates (Kieselgel 60F254, Merck; thickness 0.25 mm), eluting with a methanol-dichloromethane mixture (3-97 by volume). After elution of the zone corresponding to the

25 main product with a methanol-dichloromethane mixture (10-90 by volume) and then evaporation of the solvents under reduced pressure (0.27 kPa) at a temperature close to 40°C, 155.2 mg of 4 α ,10 β -diacetoxy-2 α -

benzoyloxy-5 β ,20-epoxy-1 β -hydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (2R,3S)-3-benzoylamino-2-hydroxy-3-phenylpropionate are obtained in the form of a white foam whose characteristics are the following:

5 - specific rotation: $[\alpha]_D^{20} = -30.5^\circ$ (c = 0.491; methanol)

- proton NMR spectrum: (300 MHz; CDCl₃; δ in ppm; coupling constants J in Hz): 1.27 (s, 3H: -CH₃, 16 or 17); 1.30 (s, 3H: -CH₃, 16 or 17); 1.40 (mt, 1H: -H7); 1.62 and 2.25 (q and m, 1H each: CH₂- of cyclopropane); 1.85 (s, 3H: -CH₃, 18); 1.96 (s, 1H: -OH in 1); 2.05 and 2.48 (d and m, 1H each: -CH₂- in 6); 2.24 (s, 3H: -COCH₃, in 10); 2.28 and 2.50 (m, 1H each: -CH₂ in 14); 2.45 (s, 3H: -COCH₃, in 4); 3.52 (d, 1H: -OH in 2'); 4.10 and 4.35 (d, 1H each: -CH₂ in 20); 4.11 (d, 1H: -H3); 4.77 (broad d, 1H: -H5); 4.82 (dd, 1H: -H2'); 5.70 (d, 1H: -H in 2); 5.84 (dd, 1H: -H3'); 6.30 (broad t, 1H: -H13); 6.36 (s, 1H: -H10); 7.00 (d, 1H: -CONH-); from 7.35 to 8.30 (m, 15H: -C₆H₅ in 3', -OCOC₆H₅ and NHCOC₆H₅).

20 The 4 α ,10 β -diacetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β -hydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (2R, 3S)-3-amino-2-hydroxy-3-phenylpropionate can be prepared by carrying out the procedure under the conditions described in Example 1 for the preparation of 4-acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,10 β -dihydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (2R,3S)-3-amino-2-hydroxy-3-phenylpropionate. Thus, starting with 1.6 g of 4 α ,10 β -diacetoxy-2 α -benzoyloxy-

5 β ,20-epoxy-1 β -hydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidincarboxylate, 1.14 g of 4 α ,10 β -diacetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β -hydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl are obtained in the form of a white foam.

The 4 α ,10 β -diacetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β -hydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidincarboxylate can be prepared under the conditions described in Example 1 for the preparation of 4 α -acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,10 β -dihydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinecarboxylate. Thus, starting with 2.2 g of 4 α ,10 β -diacetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β -hydroxy-9-oxo-7 β -trifluoromethanesulphonate-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidincarboxylate, 1.62 g of 4 α ,10 β -diacetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β -hydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidincarboxylate are obtained in the form of a white foam.

25 The 4 α ,10 β -diacetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β -hydroxy-9-oxo-7 β -trifluoromethanesulphonate-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidincarboxylate can be

prepared under the conditions described in Example 1 for the preparation of 4 α -acetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,10 β -dihydroxy-9-oxo-7 β -trifluoromethanesulphonate-19-nor-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidin-5 carboxylate. Thus, starting with 2.4 g of 4 α ,10 β -diacetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,7 β -dihydroxy-9-oxo-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinocarboxylate, 2.46 g of 10 4 α ,10 β -diacetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β -hydroxy-9-oxo-7 β -trifluoromethanesulphonate-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinocarboxylate are obtained in the form of a white foam.

15 The 4 α ,10 β -diacetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β ,7 β -dihydroxy-9-oxo-11-taxen-13 α -yl (4S,5R)-3-tert-butoxycarbonyl-2,2-dimethyl-4-phenyl-5-oxazolidinocarboxylate can be prepared under the conditions described in International Application 20 PCT WO 9209589.

EXAMPLE 3

To a solution of 550 mg of 4 α ,10 β -diacetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β -hydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (2R,3S)-3-amino-2-hydroxy-3-25 phenylpropionate in 1 cm³ of dichloromethane, kept under an argon atmosphere, are added 76 mg of sodium hydrogen carbonate and then, dropwise, at a temperature close to 20°C, a solution of 197 mg of di-tert-butyl dicarbonate

in 1 cm³ of dichloromethane. The solution obtained is stirred for 15 hours at a temperature close to 20°C and then supplemented with a mixture of 5 cm³ of distilled water and 10 cm³ of dichloromethane. The aqueous phase
5 is extracted with 5 cm³ of dichloromethane. The combined organic phases are dried over magnesium sulphate, filtered and then concentrated to dryness under reduced pressure (2.7 kPa) at 40°C. 780 mg of a white foam are thus obtained which are purified by chromatography at
10 atmospheric pressure on 50 g of silica (0.063-0.2 mm) contained in a column 2.5 cm in diameter, eluting with a methanol-dichloromethane mixture (1-99 then 2.5-97.5 by volume) and collecting 10 cm³ fractions. The fractions containing only the desired product are
15 pooled and concentrated to dryness under reduced pressure (2.7 kPa) at 40°C. 660 mg of a white foam are thus obtained. A sample of 300 mg is purified by preparative chromatography on 12 thin-layer silica plates (Kieselgel 60F254, Merck; thickness 0.25 mm),
20 eluting with a methanol-dichloromethane mixture (4-96 by volume). After elution of the zone corresponding to the main product with a methanol-dichloromethane mixture (10-90 by volume) and then evaporation of the solvents under reduced pressure (0.27 kPa) at a
25 temperature close to 40°C, 159.7 mg of 4 α ,10 β -diacetoxy-2 α -benzoyloxy-5 β ,20-epoxy-1 β -hydroxy-7 β ,8 β -methylene-9-oxo-19-nor-11-taxen-13 α -yl (2R,3S)-3-tert-butoxycarbonylamino-2-hydroxy-3-phenylpropionate are

41

obtained in the form of a white foam whose characteristics are the following:

- specific rotation: $[\alpha]_D^{20} = -34^\circ$ ($c = 0.564$; methanol)
- proton NMR spectrum: (400 MHz; CDCl_3 ; δ in ppm;
5 coupling constants J in Hz): 1.28 (s, 3H: $-\text{CH}_3$ 16 or 17); 1.30 [s, 9H: $-\text{C}(\text{CH}_3)_3$]; 1.38 (mt, 1H: $-\text{H}_7$); 1.60 (s, 3H: $-\text{CH}_3$ 16 or 17); 1.68 and 2.25 (t and m, 1H each; CH_2 - of cyclopropane); 1.85 (s, 3H: $-\text{CH}_3$ 18); 2.10 and 2.45 (d and td, 1H each: $-\text{CH}_2$ - in 6); 2.23 (s, 3H: $-\text{COCH}_3$ in 10); 2.22 and 2.40 (m, 1H each: $-\text{CH}_2$ - in 14); 2.40 (s, 3H: $-\text{COCH}_3$ in 4); 3.28 (d, 1H: $-\text{OH}$ in 2'); 4.05 and 4.22 (d, 1H each: $-\text{CH}_2$ - in 20); 4.10 (d, 1H: $-\text{H}_3$); 4.62 (broad s, 1H: $-\text{H}_2'$); 4.73 (d, 1H: $-\text{H}_5$); 5.29 (broad d, 1H: $-\text{H}_3'$); 5.37 (d, 1H: $-\text{CONH}-$); 5.67 (d, 1H: $-\text{H}$ in 2); 6.28 (broad t, 1H: $-\text{H}_{13}$); 6.33 (s, 1H: $-\text{H}_{10}$); from 7.30 to 7.45 (mt, 5H: $-\text{C}_6\text{H}_5$ in 3'); 7.51 [t, 2H: $-\text{OCOC}_6\text{H}_5$ ($-\text{H}_3$ and $-\text{H}_5$)]; 7.61 [t, 1H: $-\text{OCOC}_6\text{H}_5$ ($-\text{H}_4$)]; 8.17 [d, 2H: $-\text{OCOC}_6\text{H}_5$ ($-\text{H}_2$ and $-\text{H}_6$)].
10
15

EXAMPLE 4

- 20 To a solution of 100 mg of 10-deacetyl-baccatin III in a mixture of 3 cm^3 of tetrahydrofuran and 0.05 cm^3 of pyridine cooled to a temperature close to -78°C and kept under an argon atmosphere, is added, dropwise, 0.09 cm^3 of trifluoromethanesulphonic anhydride. The temperature is allowed
25 to rise slowly to a temperature close to 0°C over approximately one hour, then up to a temperature close to 20°C over approximately one hour. After 2 hours at a

temperature close to 20°C, 200 mg of tetrabutyl-
ammonium iodide are added, then the solution is heated
at the boiling temperature of the solvent for 15 hours.
After cooling to a temperature close to 20°C, 10 cm³ of
5 ethyl acetate and then 1 cm³ of distilled water are
added. After separation after settling has taken place,
the organic phase is dried over magnesium sulphate,
filtered and concentrated to dryness under reduced
pressure (2.7 kPa) at 40°C. 116 mg of a yellow oil are
10 thus obtained which are purified by chromatography at
atmospheric pressure on 30 g of silica (0.063-0.2 mm)
contained in a column 2.5 cm in diameter, eluting with
an ethyl acetate-dichloromethane mixture, with an
elution gradient from 0-100 to 80-20 by volume. The
15 fractions containing the desired product are pooled and
concentrated to dryness under reduced pressure
(0.27 kPa) at 40°C. 10.3 mg of 10-deacetyl-7 β ,8 β -
methylene-19-norbaccatin III are thus obtained in the
form of a white foam whose characteristics are the
20 following:

- proton NMR spectrum: (400 MHz; CDCl₃; δ in ppm;
coupling constants J in Hz): 1.14 (s, 3H: -CH₃ in 16 or
17); 1.42 (mt, 1H: -H in 7); 1.76 and 2.31 (t and m, 1H
each; CH₂ of cyclopropane); 2.07 (s, 3H: -CH₃ in 18);
25 2.15 and 2.50 (broad d and td, 1H each: CH₂- in 6); 2.30
(s, 3H: -COCH₃ in 4); 2.28 and 2.35 (m, 1H each: -CH₂ in
14); 4.11 and 4.37 (d, 1H each: -CH₂ in 20); 4.28 (d,
1H: -H₃ in 3); 4.79 (d, 1H: -H in 5); 4.88 (broad t,

43

1H: -H in 13); 5.09 (s, 1H: -H in 10); 5.66 (d, 1H: -H in 2); 7.51 [t, 2H: -OCOC₆H₅ (-H in 3 and 5)]; 7.61 [t, 1H: -OCOC₆H₅ (-H in 4)]; 8.17 [d, 2H: -OCOC₆H₅ (-H in 2 and 6)].

5 ¹³C NMR spectrum: (100 MHz; CDCl₃; δ in ppm; uncoupled; s = singlet, d = doublet; t = triplet; q = quadruplet):
15 (q, C18); 16.5 (t, C19); 20 and 27 (q, C16 and C17);
22.5 (q, -COCH₃); 26.5 (t, C6); 33 (d, C7); 35 (s, C8);
39 (d, C3); 39.5 (t, C14); 43 (s, C15); 68 (d, C13); 76
10 (t, C20); 76.2 (d, C10); 79.5 (s, C1); 80 (s, C4); 81
(d, C2); 85 (d, C5); 129 (d, C2: -OCOC₆H₅); 130 (s, C1
of -OCOC₆H₅); 130.5 (d, C3 of -OCOC₆H₅); 134 (d, C4 of
-OCOC₆H₅); 136 (s, C11); 143 (s, C12); 168 (s, -OCOC₆H₅);
171 (s, -COCH₃); 210 (s, C9).

15 The new products of general formula (I)
manifest a significant inhibitory activity with respect
to abnormal cell proliferation and possess therapeutic
properties which permit the treatment of patients
having pathological conditions associated with abnormal
20 cell proliferation. The pathological conditions include
the abnormal cell proliferation of malignant or
nonmalignant cells of various tissues and/or organs,
comprising, with no limitation being implied, muscle,
bone or connective tissues, the skin, brain, lungs, sex
25 organs, the lymphatic or renal systems, mammary or
blood cells, liver, the digestive tract, pancreas and
thyroid or adrenal glands. These pathological
conditions can also include psoriasis, solid tumours,

cancers of the ovary, breast, brain, prostate, colon, stomach, kidney or testicles, Kaposi's sarcoma, cholangioma, chorioma, neuroblastoma, Wilms' tumour, Hodgkin's disease, melanomas, multiple myelomas, lymphatic leukaemias and acute or chronic granulocytic lymphomas. The new products according to the invention are particularly useful for the treatment of cancer of the ovary. The products according to the invention can be used to prevent or retard the appearance or reappearance of the pathological conditions or to treat these pathological conditions.

The products according to the invention can be administered to a patient in various forms adapted to the chosen route of administration which is preferably the parenteral route. Parenteral administration comprises intravenous, intraperitoneal, intramuscular or subcutaneous administrations. Intraperitoneal or intravenous administration is more particularly preferred.

The present invention also comprises pharmaceutical compositions containing at least one product of general formula (I) in a sufficient quantity adapted to use in human or veterinary therapy. The compositions can be prepared according to the customary methods, using one or more pharmaceutically acceptable adjuvants, carriers or excipients. Suitable carriers include diluents, sterile aqueous media and various nontoxic solvents. Preferably, the compositions are

provided in the form of aqueous solutions or suspensions, of injectable solutions which may contain emulsifying agents, colorants, preservatives or stabilizers.

5 The choice of adjuvants or excipients may be determined by the solubility and the chemical properties of the product, the particular mode of administration and good pharmaceutical practice.

 For parenteral administration, aqueous or
10 nonaqueous sterile solutions or suspensions are used. For the preparation of nonaqueous solutions or suspensions, natural vegetable oils such as olive oil, sesame oil or paraffin oil or injectable organic esters such as ethyl oleate can be used. The aqueous sterile
15 solutions may consist of a solution of a pharmaceutically acceptable salt in solution in water. The aqueous solutions are suitable for intravenous administration in so far as the pH is appropriately adjusted and isotonicity is achieved, for example, with
20 a sufficient quantity of sodium chloride or glucose. The sterilization can be performed by heating or by any other means which does not adversely affect the composition.

 It is clearly understood that all the
25 products entering into the compositions according to the invention should be pure and nontoxic for the quantities used.

 The compositions may contain at least 0.01 %

of therapeutically active product. The quantity of active product in a composition is such that a suitable dosage can be prescribed. Preferably, the compositions are prepared such that a single dose contains about
5 0.01 to 1000 mg of active product for parenteral administration.

The therapeutic treatment can be performed concurrently with other therapeutic treatments including antineoplastic drugs, monoclonal antibodies,
10 immunotherapies or radiotherapies or biological response modifiers. The response modifiers include, with no limitation being implied, lymphokines and cytokines such as interleukins, interferons (α , β or δ) and TNF. Other chemotherapeutic agents which are useful
15 in the treatment of disorders caused by abnormal proliferation of cells include, with no limitation being implied, alkylating agents like nitrogen mustards such as mechlorethamine, cyclophosphamide, melphalan and chlorambucil, alkyl sulphonates such as busulfan,
20 nitrosoureas such as carmustine, lomustine, semustine and streptozocin, triazenes such as dacarbazine, antimetabolites such as folic acid analogues like methotrexate, pyrimidine analogues such as fluorouracil and cytarabine, purine analogues such as mercaptopurine
25 and thioguanine, natural products like vinca alkaloids such as vinblastine, vincristine and vindesine, epipodophyllotoxins such as etoposide and teniposide, antibiotics such as dactinomycin, daunorubicin.

doxorubicin, bleomycin, plicamycin and mitomycin,
enzymes such as L-asparaginase, various agents such as
coordination complexes of platinum like cisplatin,
substituted ureas like hydroxyurea, methylhydrazine
5 derivatives such as procarbazine, adrenocortical
suppressants such as mitotane and aminoglutethimide,
hormones and antagonists such as adrenocorticosteroids
such as prednisone, progestins such as
hydroxyprogesterone caproate, methoxyprogesterone
10 acetate and megestrol acetate, oestrogens such as
diethylstilbestrol and ethynylestradiol, antioestrogens
such as tamoxifen, and androgens such as testosterone
propionate and fluoxymesterone.

The doses used for carrying out the methods
15 according to the invention are those which permit a
prophylactic treatment or a maximum therapeutic
response. The doses vary according to the form of
administration, the particular product selected and the
characteristics specific to the subject to be treated.
20 In general, the doses are those which are
therapeutically effective for the treatment of
disorders caused by abnormal cell proliferation. The
products according to the invention can be administered
as often as necessary to obtain the desired therapeutic
25 effect. Some patients may respond rapidly to relatively
high or low doses, and then require low or zero
maintenance doses. Generally, low doses will be used at
the beginning of the treatment and, if necessary,

increasingly higher doses will be administered until an optimum effect is obtained. For other patients, it may be necessary to administer maintenance doses 1 to 8 times per day, preferably 1 to 4 times according to the physiological needs of the patient considered. It is also possible that for certain patients it may be necessary to use only one to two daily administrations.

In man, the doses are generally between 0.01 and 200 mg/kg. For intraperitoneal administration, the doses will generally be between 0.1 and 100 mg/kg and, preferably, between 0.5 and 50 mg/kg and, still more specifically, between 1 and 10 mg/kg. For intravenous administration, the doses are generally between 0.1 and 50 mg/kg and, preferably, between 0.1 and 5 mg/kg and, still more specifically, between 1 and 2 mg/kg. It is understood that, in order to choose the most appropriate dosage, account should be taken of the route of administration, the patient's weight, his general state of health, his age and all factors which may influence the efficacy of the treatment.

The following example illustrates a composition according to the invention.

EXAMPLE

40 mg of the product obtained in Example 1 are dissolved in 1 cm³ of Emulphor EL 620 and 1 cm³ of ethanol and then the solution is diluted by addition of 18 cm³ of physiological saline.

The composition is administered by perfusion

for 1 hour by introduction into physiological saline.

5 Although the invention has been described in conjunction
with specific embodiments, it is evident that many alternatives
and variations will be apparent to those skilled in the art in
light of the foregoing description. Accordingly, the invention
is intended to embrace all of the alternatives and variations
10 that fall within the spirit and scope of the appended claims.
The above references are hereby incorporated by reference.